

Indication		Grp.no. Medicament	
		Group 3. Pain (Continued)	
Severe pains Major wounds Injuries Burns Fractures pre- and post operative pains  MFAG tables 7,8,10,13  <b>A, B1,B2, C, P</b>	3.4	<b>Medicinal substance:</b>	Nubain (Nalbufine)
		<b>Form:</b>	Nubain: Nalbufine, ampoule 10mg/1ml, 10 x 1 ml single dose
		<b>Effect:</b>	Analgesic. Takes effect after 2- 3 min. Effective for 3 - 6 hours.
		<b>Dosage:</b>	Inject sc or im: effect within 15 min. Inject iv: effect within 2 to 3 min. 10 - 20 mg , if needed every 3 - 6 hours
		<b>Side-effects:</b>	Sedation, headache, nausea and vomiting.
		<b>Validity:</b>	3 years
		<b>Storage:</b>	Room temperature ( 15 - 25 degrees C)
		<i>Remarks</i>	Nalbufine is related to morphine and the effects are comparable. Do not use alcohol, sleeping medicine or other medicines that cause drowsiness at the same time. In case of an overdose a patient may stop breathing. Antidote: Naloxone. Consult RADIO MEDICAL.

		Gruppe 3. Smertestillende midler (Continued)	
Sterke smerter Større sår Skader Brandsår Brud Pre- og postoperative smerter  MFAG tabeller 7,8,10,13  <b>A, B1,B2, C, P</b>	3.4	<b>Lægemiddelstof:</b>	Nubain (Nalbufine)
		<b>Form:</b>	Nubain: Nalbufine ampuller 10 mg/1 ml, 10 x 1ml
		<b>Virkning:</b>	Smertestillende. Virkningen indtræder efter 2 - 3 min. og varer 3 - 6 timer.
		<b>Dosering:</b>	Injektion s.c./i.m.: virkning efter 15 min. Injektion i.v.: virkning efter 2 - 3 min. 10 - 20 mg, efter behov hver 3 - 6 timer
		<b>Bivirkninger:</b>	Sløvhed, hovedpine, kvalme og opkastning.
		<b>Holdbarhed:</b>	3 år
		<b>Opbevaring:</b>	Rumtemperatur (15-25°C)
		<i>Bemærkninger:</i>	Nalbufine er beslægtet med morfin og besidder de samme virkninger. Brug ikke alkohol, sovetabletter eller andre lægemidler som samtidig forårsager dødsghed. Ved overdosering kan patientens åndedæt stoppe. Modgift: Naloxone. Søg samråd med RADIO MEDICAL.