

## Indication

## Grp.no. Medicament

|                 |     | <b>Group 3. Pain</b>        |  | <b>(Continued)</b> |
|-----------------|-----|-----------------------------|--|--------------------|
| Severe pains    | 3.3 | <b>Medicinal substance:</b> | Pentazocine  |                    |
|                 |     | <b>Form:</b>                | Tablets of 50 mg   |                    |
|                 |     | <b>Effect:</b>              | Analgesic. Takes effect after 15-30 min. Effective for at least 3 hours.   |                    |
|                 |     | <b>Dosage:</b>              | 1 - 2 tablets every 3 - 4 hours. Maximum 12 tablets per day  |                    |
|                 |     | <b>Side-effects:</b>        | Drowsiness, nausea, vomiting, dizziness, headache, obstipation.  |                    |
|                 |     | <b>Validity:</b>            | 3 years  |                    |
|                 |     | <b>Storage:</b>             |  |                    |
|                 |     | <i>Remarks</i>              | Pentazocine is related to morphine and the effects are comparable. Do not use alcohol, sleeping medicine or other medicines that cause drowsiness at the same time. Consult RADIO MEDICAL. |                    |
| <b>A, B1,B2</b> |     |                             |  |                    |

|                 |     | <b>Gruppe 3. Smertestillende midler</b> |  | <b>(Continued)</b> |
|-----------------|-----|---|--|--------------------|
| Sterke smerter  | 3.3 | <b>Lægemedelstof:</b>                   | Pentazocine  |                    |
|                 |     | <b>Form:</b>                            | Tabletter 50 mg  |                    |
|                 |     | <b>Virkning:</b>                        | Smertestillende. Virkningen indtræder efter 15 - 30 min. og varer mindst 3 timer.  |                    |
|                 |     | <b>Dosering:</b>                        | 1 - 2 tabletter hver 3 - 4 time. Maksimum 12 tabletter per dag.  |                    |
|                 |     | <b>Bivirkninger:</b>                    | Døsighed, kvalme, opkastning, svimmelhed, hovedpine, forstoppelse.   |                    |
|                 |     | <b>Holdbarhed:</b>                      | 3 år   |                    |
|                 |     | <b>Opbevaring:</b>                      |  |                    |
|                 |     | <i>Bemærkninger:</i>                    | Pentazocine er beslægtet med morfin og besidder de samme virkninger. Brug ikke alkohol, sovetabletter eller andre lægemidler som samtidig forårsager døsighed. Søg samråd med RADIO MEDICAL. |                    |
| <b>A, B1,B2</b> |     |   |  |                    |